

# A structure factor approach to the Sunyaev-Zel'dovich effect

Alfredo Sandoval-Villalazo<sup>\*†</sup>

<sup>\*</sup>*Relativity and Cosmology Group, School of Computer Science and Mathematics, Portsmouth University,  
Portsmouth PO1 2EG, United Kingdom*

<sup>†</sup>*Departamento de Ciencias, Universidad Iberoamericana, Col. Lomas de Santa Fe 01210, Mexico D.F., Mexico*

Structure factor theory (SFT) is applied to analyze distortions to the Cosmic Microwave Background Radiation (CMBR) due the Sunyaev-Zel'dovich Effect (SZE) in typical clusters of galaxies. It is shown that a suitable structure factor associated to a collisionless (Knudsen) gas is able to reproduce the SZE distortion of the CMBR. Physical arguments suggest that SFT yields an approximate solution to the Kompaneets equation and sets plausible values for the two “free” parameters involved in the calculation of the spectral distortion. Finally, further applications of the SFT to other standard cosmological problems are briefly discussed.

## I. Introduction

For decades, light scattering theory has been applied to the analysis of time-dependent fluctuations of the thermodynamical variables of simple fluids. It is a well-known fact that the Laplace-Fourier transform of the time correlation function of density fluctuations gives rise to a *scattering law* that describes the general behavior of the scattered spectra. The mathematical framework in which the scattering law is established in different cases is called Structure Factor Theory (SFT). Structure factors have been studied from different points of view, mainly involving kinetic theory [1] and hydrodynamics [2]. For astrophysical systems, low densities prevent the calculation of structure factors in terms of hydrodynamical models, but the general form of the appropriate scattering law for a collisionless (Knudsen) gas is perfectly established [3]. Thus, the application of SFT to the analysis of distortions of the Cosmic Microwave Background Radiation (CMBR) is possible. Specifically we shall concentrate in CMBR scattering by the hot gas present in clusters of gases, better known as the thermal Sunyaev-Zel'dovich Effect (SZE).

The SZE [5] provides a potentially very important probe of large-scale structure and the main cosmological parameters (see e.g. [6,7]). The SZE is associated to Compton scattering of CMBR photons by hot electrons in the ionized intra-cluster gas. On average, photons are shifted to higher frequencies, but the effect vanishes at a critical frequency where non-relativistic calculations give  $\sim 217$  GHz for typical gas temperatures  $\sim 10^8$  K (Relativistic corrections to the SZE [8] introduce only a very small increase in the critical frequency). The first approach to the SZE was made by means of the Kompaneets diffusion equation, which describes photon diffusion in an electron gas.

This paper shows that the CMBR distortions associated with the thermal SZE can be reproduced by means of SFT. The picture is very simple. As Rephaeli pointed out [8], most photons are not scattered *even once* by the gas components. Thus, if we consider a single frequency in the CMBR photons, the corresponding line will be slightly broadened due to the scattering of some photons. This type of broadening can be described by SFT. The whole spectral distortion is then described by convolution-type integrals that have already been applied in other CMBR problems [9]. The resulting curves describing spectral distortion will represent solutions to the Kompaneets equation. The underlying physical basis of these solutions deserves further attention.

This paper is divided as follows: section 2 is dedicated to the analysis of the SFT approach to the thermal SZE in terms of the scattering law suggested in Ref. [3]. Section 3 includes a critical comparison of the curve obtained in section 2 and the SZE curve obtained by means of the Kompaneets equation. Finally, ideas related to the physical interpretation of the “free” parameters involved in the structure factor description of the SZE, and further applications of this approach, close this work.

## II. Scattering law for a simple collisionless gases.

A very simple way of deriving the scattering law for a collisionless gas [3] starts by considering the generic form of a Maxwell-Boltzmann distribution function of speeds for a simple, non-relativistic gas.

$$f(v) = A v^2 e^{-\eta(v)^2} \quad (1)$$

The most probable velocity for a given  $\eta$  is:

$$v_o^2 = \frac{1}{\eta} \quad (2)$$

The probability that a particle has a velocity  $\mathbf{v}$  is proportional to  $e^{-\left(\frac{v}{v_o}\right)^2}$ . The displacement of a particle in time  $\tau$  through the distance  $\mathbf{r}$  corresponds to  $\mathbf{v} \tau$ . Hence, the normalized probability of finding it at  $\mathbf{r}$  at time  $\tau$ , after it was at  $(0, 0)$ , is

$$G = \frac{1}{\pi^{\frac{3}{2}} |v_o \tau|^3} e^{-\frac{r^2}{(v_o \tau)^2}} \quad (3)$$

Performing a Fourier transform in space and a Laplace transform in time of Eq. (3), the scattering law for the gas corresponds to the *structure factor*:

$$S(\nu) = \frac{1}{\sqrt{\pi} k v_o} e^{-\frac{(\nu)^2}{(k v_o)^2}} = \frac{1}{\sqrt{\pi} \alpha} e^{-\frac{(\nu)^2}{\alpha^2}} \quad (4)$$

In a collisionless gas, particles are not interacting. It is known that, in this case, the intensity of radiation scattered off the fluid will be changed from its initial value by an amount proportional to the Laplace-Fourier transform of the density time correlation function, i.e. Eq. (4) [10]. Physically, Eq.(4) represents a broadening of a single spectral line of frequency  $\nu$  due to the scattering of incoming photons off the single particles of the collisionless gas. A simple thumb rule corresponding to this broadening is:

$$\delta(\nu) \longrightarrow \frac{1}{\sqrt{\pi} \alpha} e^{-\frac{(\nu)^2}{\alpha^2}} = S(k, \nu) \quad (5)$$

Now, if the intensity of the incoming radiation, per unit of solid angle has a blackbody spectrum:

$$I_o(\nu) = \frac{2h\nu^3}{c} \frac{1}{e^{\frac{h\nu}{k_B T}} - 1} \quad (6)$$

then, the full distorted spectrum will be given by the convolution-type integral:

$$I(\nu) = \int_0^\infty I_o(\omega) S(k, \omega - \nu) d\omega \quad (7)$$

and the departures from the perfect black-body spectrum will be given by:

$$\Delta I(\nu) = I(\nu) - I_o(\nu) \quad (8)$$

In order to show how  $\Delta I(\nu)$  behaves, let us suppose that CMBR photons are traveling across a very hot gas,  $T = 10^8 K$ . The typical broadening parameter  $\alpha$  can be estimated for low density systems (as the hot intra-cluster gas of the SZE) by:

$$\alpha \sim \frac{2\pi}{c} \sqrt{\frac{k_b T}{m}} \nu \quad (9)$$

Where  $k_b$  is Boltzmann's constant and  $m$  is a single particle mass. Fig. 1 shows  $\Delta I(\nu)$  taking  $m = 1.6 \times 10^{-24}$  g (the proton mass). As a reference, we plot in Fig. 1 the SZE curve for the same temperature with Compton parameter  $y = 10^{-4}$ , as in [12]. More information about this parameter will be given in the next section. Not only the shape of the curves are very much alike, but also the order of magnitude for the CMBR distortion is practically the same.




FIG. 1. The thermal SZE standard curve (solid) compared to the CMBR spectrum distortion computed from SFT (dashed),  $\Delta I$  ( $\text{erg s}^{-1} \text{Hz}^{-1} \text{cm}^{-2} \text{ster}^{-1} \times 10^{-18}$ ), against frequency ( $\text{Hz}$ ) for  $T = 10^8 \text{K}$ .

### III. The $y$ Compton parameter in SFT

Up to this point, it has been shown that, mathematically, SZE-type curves can be obtained by means of the convolution integrals (7). Physically, the  $\alpha$  parameter is a simple line broadening proportional to the width parameter of the low density system. Of course, no sound waves exist in a system such as the hot intra-cluster gas (densities are close to  $10^{-3}$  particles per cubic centimeter), but the broadening does exist and has been the subject of several studies regarding the transition from a collision-dominated system, to a collisionless one [1]. In dense systems, sound waves will also scatter off the CMBR photons and the structure factor would have to take into account Brillouin scattering [2]. The thumb rule (5) would then involve Lorentzian shapes as in the case of liquids [10].

Nevertheless, more physics is needed to get a better description of the SZE curves. In particular, the scattering cross section between photons and particles, and the density and size of the cluster enter in the ordinary approach to the SZE by means of the *Compton “y” parameter*:

$$y = \int n \sigma \frac{m_e c^2}{k_b T} d\ell \quad (10)$$

Here  $n$  is the number density of the electrons in the cluster,  $\sigma$  is the Thomson cross section and the integral is supposed to be taken along the line of sight of a photon. Roughly, we can write:

$$y \sim n \sigma \frac{m_e c^2}{k_b T} \ell \quad (11)$$

In this case,  $m_e$  is the electron mass. For typical temperatures ( $T = 10^8 \text{K}$ ):

$$y \sim 10^{-4} \quad (12)$$

The actual value for  $y$  depends on the precise properties of the cluster. For the rest of the paper we will use  $y \sim 10^{-4}$  as a reference value [12]. In the ordinary approach to the SZE,  $\Delta I(\nu) = y f(\nu, T)$ , where  $f(\nu, T)$  is a spectral distortion function usually computed by means of an approximate solution of the Kompaneets diffusion equation. The precise form of  $f(\nu, T)$  is not important for the present paper and the reader is encouraged to consult the reviews. One interesting fact is that the change of brightness temperature of the CMBR in the Rayleigh-Jeans region is simply given by:

$$\frac{\Delta T_{CMB}}{T_{CMB}} = -2y \quad (13)$$

Now, the  $y$  factor may enter into the SFT approach to the SZE if we slightly modify Eq. (4) so that the distorted spectrum will now read:

$$I(\nu) = \int_0^\infty I_o(\omega) S(\omega - (1 - 2y)\nu) d\omega \quad (14)$$

Fig. 2 shows that the “fine tuning” given by the  $y$  parameter is very effective. Indeed, all the features of the SZE are reproduced accurately enough to consider SFT as an appropriate tool to analyze CMBR distortions.




FIG. 2. The same as in Fig. 1, but taking into account Eq. (14). The reproduction of the main features of the SZE curve is clear.

#### IV. Discussion

There is a physical reason that supports the crucial center shift in the convolution integral (14). Compton effect generates a shift in frequency that, integrated along the photon path in the cluster, roughly indicates that the center of the structure factor, for a fixed frequency  $\nu$ , must be displaced by an amount proportional to the “ $y$ ” parameter. This picture is consistent with the description of comptonization in clusters included in Rybicki and Lightman’s text, in which the “ $y$ ” parameter is defined as a tool to determine whether a photon will significantly change its energy (frequency) while traversing the medium [11]. Thus, “ $y$ ” is basically the average fractional frequency change due to Compton scattering, considering all its flight across the intra-cluster gas. The calculation here presented *is not* a simple fit. Moreover, if the Compton factor is reduced by a factor of 10 (corresponding to a *physical* multiplying factor of 0.325 for  $\alpha$  in Eq.(9)), the reproduction of the SZE curve by means of Eq.(14) is practically perfect.




FIG. 3. The same as in Fig. 2, but with the Compton factor lowered by a factor of 10

All parameters included in the basic equation (14) have a proper physical meaning and other uses of the formalism in CMBR related problems seem plausible. To the author's knowledge, this is the first time that SZE-type curves are obtained by means of convolution integrals. The basic curve can be associated with a standard structure factor, while "fine tuning" can be related to the gas comptonization parameter " $y$ ". Obviously, the integral expression (14) must be related to the standard solution of the Kompaneets equation that ordinarily describes the non-relativistic SZE.

The work here presented has established some basics about a SFT approach to the SZE. The results shown here suggest that structure factors may provide alternative approaches to CMBR distortion problems. Since line broadening is also present in all dissipative systems, proper convolution integrals involving Lorentzian structure factors may yield alternative approaches to light scattering in cosmological processes, such as recombination. In recombination, photon scattering due to acoustic waves must be taken into account. Other future work includes the relativistic generalization of this work in terms of a proper Maxwellian. Finally, a kinetic theory approach to structure factor theory by means of the techniques suggested in Ref. [14] will soon be addressed.

The author thanks Leopoldo S. García-Colín (UAM-I) and Roy Maartens (Portsmouth) for valuable comments and suggestions. This work was supported by a CONACyT postdoctoral grant. The author also thanks the Relativity and Cosmology Group at Portsmouth for hosting him while this work was done.

- 
- [1] S. Yip and M. Nelkin, Phys. Rev. 135A, 1241 (1964).
  - [2] B.J. Berne and R. Pecora, *Dynamic Light Scattering* (Wiley, N.Y., 1976).
  - [3] P.A. Egelstaff, *An Introduction to the Liquid State* (Academic Press, London and New York, 1967).
  - [4] H.E. Stanley, *Introduction to phase transitions and critical phenomena*, (Oxford University Press, Oxford, 1971).
  - [5] Y.B. Zel'dovich and R.A. Sunyaev, Astrophys. Space Sci. **4**, 30 (1969); R.A. Sunyaev and Y.B. Zel'dovich, Comments Astrophys. Space Phys. **4**, 173 (1972).
  - [6] M. Birkinshaw, Phys. Rep. **310**, 97 (1999).
  - [7] J.M. Diego, E. Martinez-Gonzalez, J.L. Sanz, L. Benitez and J. Silk, Mon. Not. R. Astron. Soc., to appear (astro-ph/0103512).
  - [8] Y. Rephaeli, Astrophys. J. **445**, 33 (1995); A.D. Challinor and A.N. Lasenby, Astrophys. J. **499**, 1 (1998); *ibid.*, Astrophys. J. **510**, 930 (1999); S.Y. Sazonov and R.A. Sunyaev, Astrophys. J. **508**, 1 (1998).
  - [9] R. Di Stefano, L.H. Ford, H. Yu and D.J. Fixsen (2001) astro-ph/0107001. Submitted to Astrophys. J.
  - [10] B.J. Berne and R. Pecora, *Dynamic Light Scattering* (Wiley, N.Y., 1976).
  - [11] G.B. Rybicki and A.P. Lightman *Radiative Processes in Astrophysics* (Wiley, N.Y., 1979).

- [12] J.E. Carlstrom, M. Joy, L. Gregol G. Holder, W.L.Holzapfel, S. LaRoque, J.J. Mohr and E.D. Reese (astro-ph/0103480), to appear in *Constructing the Universe with Clusters of Galaxies*, IAP conference (2000).
- [13] Y. Rephaeli, (2001) astro-ph/0105192. Submitted to world Scientific.
- [14] A. Sandoval-Villalbazo and L.S. Garcia-Colin, Physica A 278 (2000) 428.